

# Validation Report

## Rapid Test Easy for Gluten

Cat.# M2243

**The quick detection for wheat proteins (gluten)  
in swabs and clean-in-place rinse water**

**Detection Limit            0.5 µg/mL wheat protein  
                                     0.425 µg/mL gluten**

**Morinaga Institute of Biological Science, Inc.**

**Sachiura 2-1-16, Kanazawa-Ku,  
Yokohama-Shi,  
236-0003, Japan**

**TEL: +81-45-791-7674  
FAX: +81-45-791-7675**

**E-mail for inquiries: [info\\_miobs\\_e@morinaga.co.jp](mailto:info_miobs_e@morinaga.co.jp)**

## 1. Scope

The Rapid Test Easy for Gluten is a lateral flow method for quick determination of whether there are residual wheat proteins (gluten) of allergenic ingredients in swabs and Clean-in-place rinse water. Water or phosphate buffered-saline(PBS(-)) can be used to prepare test solutions.

## 2. Overview

Detection protein	Gliadin
Detection limit	0.5 µg/mL wheat protein 0.425 µg/mL gluten
Reaction time	10 min

## 3. Sensitivity

Wheat solution protein concentration	in water	in PBS(-)
0 ppm	-	-
0.5 ppm	+	+
20 ppm	+	+
200 ppm	+	+

+: Positive    ±: Weak positive  
-: Negative

\*Protein concentration was valued using a 2-D Quant Kit (GE Health care)

Boiled wheat solution protein concentration	in water	in PBS(-)
0 ppm	-	-
0.5 ppm	+	+
20 ppm	+	+
200 ppm	+	-

\*Protein concentration was valued using a 2-D Quant Kit (GE Health care)

## 4. Confirmation of Prozone Phenomenon

Material	in water	in PBS(-)
0.1% Wheat	+	+

## 5. Cross-Reactivity

Material	0.1 % material in water	0.1 % material in PBS(-)
Egg	-	-
Milk	-	-
Buckwheat	-	-
Soy bean	-	-
Corn flour	-	-
Peanut (Roasted)	-	-
Rice flour	-	-
Corn flour	-	-
Shrimp	-	-
Crab	-	-